

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MUC1 / EMA / CD227 (Epithelial Marker) (rMUC1/960), Biotin conjugate, 0.1mg/mL, Clone: [rMUC1/960], Mouse, Monoclonal BOT-BNCB0960-100**

Artikelname	MUC1 / EMA / CD227 (Epithelial Marker) (rMUC1/960), Biotin conjugate, 0.1mg/mL, Clone: [rMUC1/960], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0960-100
Hersteller Artikelnummer	BNCB0960-100
Alternativnummer	BOT-BNCB0960-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Human milk-fat globule membranes (HMFGM)
Konjugation	Biotin
Produktbeschreibung	This MAb reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. The dominant epitope of this MAb involves both amino acids as well as sugar moieties. Neuraminidase treatment destroys the an...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rMUC1/960]

Molekulargewicht	265-400 kDa
UniProt	<a href="#">P15941</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user