

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Ep-CAM / CD326(Cocktail PAN-EpCAM), Biotin conjugate, 0.1mg/mL, Clone: [Cocktail PAN-EpCAM], Mouse, Monoclonal BOT-BNCB0969-500**

Artikelname	Ep-CAM / CD326(Cocktail PAN-EpCAM), Biotin conjugate, 0.1mg/mL, Clone: [Cocktail PAN-EpCAM], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0969-500
Hersteller Artikelnummer	BNCB0969-500
Alternativnummer	BOT-BNCB0969-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	Exosome Staining, FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Ep-CAM protein (full-length and fragments)
Konjugation	Biotin
Produktbeschreibung	This is a cocktail of four highly specific monoclonal antibodies (EGP40/826, EGP40/837, EGP40/1110, EGP40/1120) that recognize extracellular as well as intracellular domains of the epithelial cellular adhesion molecule (Ep-CAM). Epithelial specific a...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[Cocktail PAN-EpCAM]

Molekulargewicht	40-43 kDa
UniProt	<a href="#">P16422</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user