

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD54(W-CAM-1, same as Wehi-CAM-1 or 1H4), Biotin conjugate, 0.1mg/mL, Clone: [W-CAM-1, same as Wehi-CAM-1 or 1H4], Mouse, Monoclonal BOT-BNCB0988-100

Artikelname	CD54(W-CAM-1, same as Wehi-CAM-1 or 1H4), Biotin conjugate, 0.1mg/mL, Clone: [W-CAM-1, same as Wehi-CAM-1 or 1H4], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0988-100
Hersteller Artikelnummer	BNCB0988-100
Alternativnummer	BOT-BNCB0988-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Raji Burkitt lymphoma cells
Konjugation	Biotin
Produktbeschreibung	Recognizes an 85-115 kDa protein (variation with cell type), identified as intercellular adhesion molecule (ICAM-1) (Workshop IV). It has 7 potential N-linked glycosylation sites. ICAM-1 is a single chain glycoprotein of Ig supergene family, present ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[W-CAM-1, same as Wehi-CAM-1 or 1H4]
Molekulargewicht	85-115 kDa
UniProt	P05362
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Functional Studies order Ab without BSA & azide) Optimal dilution for a specific application should be determined by user