

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytochrome C(CYCS/1010), Biotin conjugate, 0.1mg/mL, Clone: [CYCS/1010], Mouse, Monoclonal BOT-BNCB1010-500

| | |
|--------------------------|---|
| Artikelname | Cytochrome C(CYCS/1010), Biotin conjugate, 0.1mg/mL, Clone: [CYCS/1010], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1010-500 |
| Hersteller Artikelnummer | BNCB1010-500 |
| Alternativnummer | BOT-BNCB1010-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC, WB |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant full-length human CYCS protein |
| Konjugation | Biotin |
| Produktbeschreibung | Cytochrome c is a well-characterized mobile electron transport protein that is essential to energy conversion in all aerobic organisms. In mammalian cells, this highly conserved protein is normally localized to the mitochondrial inter-membrane space.... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CYCS/1010] |

| | |
|------------------------|--|
| Molekulargewicht | 15 kDa |
| UniProt | P99999 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |