

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD47(IAP/964 + B6H12.2), Biotin conjugate, 0.1mg/mL, Clone: [IAP/964 B6H12.2], Mouse, Monoclonal BOT-BNCB1029-500

| | |
|--------------------------|---|
| Artikelname | CD47(IAP/964 + B6H12.2), Biotin conjugate, 0.1mg/mL, Clone: [IAP/964 B6H12.2], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1029-500 |
| Hersteller Artikelnummer | BNCB1029-500 |
| Alternativnummer | BOT-BNCB1029-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human IAP protein (IAP/964), Intact CD47 purified from placenta (B6H12.2) |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody reacts with Ig domain of CD47 protein. CD47, originally named integrin-associated protein (IAP), is a 50 kDa protein containing five membrane-spanning sequences and a short cytoplasmic tail. CD47 plays a role in both cell adhesion by ac... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [IAP/964 B6H12.2] |
| Molekulargewicht | ~50 kDa |
| UniProt | Q08722 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |