

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD38(AT2), Biotin conjugate, 0.1mg/mL, Clone: [AT2], Mouse, Monoclonal BOT-BNCB1086-500

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | CD38(AT2), Biotin conjugate, 0.1mg/mL, Clone: [AT2], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1086-500 |
| Hersteller Artikelnummer | BNCB1086-500 |
| Alternativnummer | BOT-BNCB1086-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | human T cell line CCRF-CEM |
| Konjugation | Biotin |
| Produktbeschreibung | This MAb reacts with a 45 kDa glycopeptide, which is a type II membrane glycoprotein with a transmembrane sequence near the NH2 terminus. CD38 is a type II transmembrane glycoprotein that is present on early B- and T-cell lineages and activated B- an... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [AT2] |
| Molekulargewicht | ~45 kDa (Glycoprotein), 35 kDa (protein core) |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P28907 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |