

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD44 Standard(HCAM/1097), Biotin conjugate, 0.1mg/mL, Clone: [HCAM/1097], Mouse, Monoclonal BOT-BNCB1097-100

| | |
|--------------------------|---|
| Artikelname | CD44 Standard(HCAM/1097), Biotin conjugate, 0.1mg/mL, Clone: [HCAM/1097], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1097-100 |
| Hersteller Artikelnummer | BNCB1097-100 |
| Alternativnummer | BOT-BNCB1097-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human HCAM protein |
| Konjugation | Biotin |
| Produktbeschreibung | Recognizes a cell surface glycoprotein of 80-95 kDa (CD44) on lymphocytes, monocytes, and granulocytes. The CD44 family of glycoproteins exists in a number of variant isoforms, the most common being the standard 85-95 kDa or hematopoietic variant (CD... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [HCAM/1097] |
| Molekulargewicht | 80-95 kDa |
| UniProt | P16070 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |