

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cyclin B1(CCNB1/1098), Biotin conjugate, 0.1mg/mL, Clone: [CCNB1/1098], Mouse, Monoclonal BOT-BNCB1098-100

| | |
|--------------------------|---|
| Artikelname | Cyclin B1(CCNB1/1098), Biotin conjugate, 0.1mg/mL, Clone: [CCNB1/1098], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1098-100 |
| Hersteller Artikelnummer | BNCB1098-100 |
| Alternativnummer | BOT-BNCB1098-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human full-length CCNB1 protein |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a protein of 55-62 kDa, identified as cyclin B1. In mammals, cyclin B associates with inactive p34cdc2, which facilitates phosphorylation of p34cdc2 at aa 14Thr and 15Tyr. This maintains the inactive state until the end of G2... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CCNB1/1098] |

| | |
|------------------------|--|
| Molekulargewicht | 55-62 kDa |
| UniProt | P14635 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not stain methanol-fixed cells Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |