

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Arginase 1 (Hepatocellular Carcinoma Marker) (ARG1/1125) , Biotin conjugate, 0.1mg/mL, Clone: [ARG1/1125], Mouse, Monoclonal BOT-BNCB1125-100**

Artikelname	Arginase 1 (Hepatocellular Carcinoma Marker) (ARG1/1125) , Biotin conjugate, 0.1mg/mL, Clone: [ARG1/1125], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1125-100
Hersteller Artikelnummer	BNCB1125-100
Alternativnummer	BOT-BNCB1125-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human ARG1 protein fragment (around aa11-97) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a protein of 35-38 kDa, which is identified as Arginase 1 (ARG1). Arginase is a manganese metallo-enzyme that catalyzes the hydrolysis of arginine to generate ornithine and urea. Arginase I and II are isoenzymes which differ ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[ARG1/1125]
Molekulargewicht	35-38 kDa
UniProt	<a href="#">P05089</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user