

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MHC II (HLA-DP) (beta chain)(Bra-14), Biotin conjugate, 0.1mg/mL, Clone: [Bra-14], Mouse, Monoclonal BOT-BNCB1129-100

Artikelname	MHC II (HLA-DP) (beta chain)(Bra-14), Biotin conjugate, 0.1mg/mL, Clone: [Bra-14], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1129-100
Hersteller Artikelnummer	BNCB1129-100
Alternativnummer	BOT-BNCB1129-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human REH cells
Konjugation	Biotin
Produktbeschreibung	Reacts with a common epitope of human major histocompatibility (MHC) class II antigens, HLA-DR and DP. Human MHC class II antigens are transmembrane glycoproteins composed of an alpha chain (36 kDa) and a beta chain (27 kDa). They are expressed prima...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[Bra-14]
Molekulargewicht	36 kDa (? chain) and 27 kDa (? chain)
UniProt	P04440
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user