

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Complement C4d(TDM16), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [TDM16], Mouse, Monoclonal BOT-BNCB1232-500

| | |
|--------------------------|---|
| Artikelname | Complement C4d(TDM16), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [TDM16], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1232-500 |
| Hersteller Artikelnummer | BNCB1232-500 |
| Alternativnummer | BOT-BNCB1232-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Human Complement 4d protein |
| Konjugation | Biotin |
| Produktbeschreibung | This MAb is specific to Complement 4d (C4d) and it reacts with the secreted as well as cell-bound C4d.C4d is a degradation product of the activated complement factor C4b. Complement 4b is typically activated by binding of Abs to specific target molec... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [TDM16] |
| Molekulargewicht | 192 kDa (predicted) |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| UniProt | P0COL4 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |