

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **PTH (Parathyroid Hormone) (N & C Terminal) (3H9 + PTH/1175), Biotin conjugate, 0.1mg/mL, Clone: [3H9 PTH/1175], Mouse, Monoclonal BOT-BNCB1236-100**

|                          |   |
|--------------------------|---|
| Artikelname              | PTH (Parathyroid Hormone) (N & C Terminal) (3H9 + PTH/1175), Biotin conjugate, 0.1mg/mL, Clone: [3H9 PTH/1175], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNCB1236-100  |
| Hersteller Artikelnummer | BNCB1236-100  |
| Alternativnummer         | BOT-BNCB1236-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Synthetic peptide corresponding to amino acids 1 to 34 of PTH conjugated to a carrier (3H9), Recombinant fragment (84 amino acid residues from C-terminus) of human PTH protein   |
| Konjugation              | Biotin  |
| Produktbeschreibung      | The epitope of this MAb maps in the C-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood calcium levels by dissolving the salts i... |
| Klonalität               | Monoclonal  |

|                        |  |
|------------------------|--|
| Konzentration          | 0.1 mg/mL  |
| Klon-Bezeichnung       | [3H9 PTH/1175]   |
| Molekulargewicht       | 9 kDa  |
| UniProt                | <a href="#">P01270</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |