

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **p27 / KIP1(DCS-72.F6 + KIP1/769 (SX53G8)), Biotin conjugate, 0.1mg/mL, Clone: [DCS-72.F6 KIP1/769], Mouse, Monoclonal BOT-BNCB1237-500**

Artikelname	p27 / KIP1(DCS-72.F6 + KIP1/769 (SX53G8)), Biotin conjugate, 0.1mg/mL, Clone: [DCS-72.F6 KIP1/769], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1237-500
Hersteller Artikelnummer	BNCB1237-500
Alternativnummer	BOT-BNCB1237-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Monkey, Mouse, Rat
Immunogen	Mouse recombinant p27 protein (DCS-72.F6), Recombinant human CDKN1B protein (KIP1/769)
Konjugation	Biotin
Produktbeschreibung	Recognizes a 27 kDa protein, identified as the p27/Kip1, a cell cycle regulatory mitotic inhibitor. Its epitope spans between aa 83-204 of p27. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. p27/Kip1 function...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[DCS-72.F6 KIP1/769]
Molekulargewicht	25-26 kDa
UniProt	<a href="#">P46527</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user