

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD11c (Dendritic Cell Marker)(ITGAX/1242), Biotin conjugate, 0.1mg/mL, Clone: [ITGAX/1242], Mouse, Monoclonal BOT-BNCB1242-500

| | |
|--------------------------|---|
| Artikelname | CD11c (Dendritic Cell Marker)(ITGAX/1242), Biotin conjugate, 0.1mg/mL, Clone: [ITGAX/1242], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1242-500 |
| Hersteller Artikelnummer | BNCB1242-500 |
| Alternativnummer | BOT-BNCB1242-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human ITGAX protein fragment (aa 637-827) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | Recognizes a protein of 145 kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (MAC-1) and CD11... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [ITGAX/1242] |
| Molekulargewicht | 145 kDa |
| UniProt | P20702 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |