

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD11c (Dendritic Cell Marker) (ITGAX/1243) , Biotin conjugate, 0.1mg/mL, Clone: [ITGAX/1243], Mouse, Monoclonal BOT-BNCB1243-500

| | |
|--------------------------|---|
| Artikelname | CD11c (Dendritic Cell Marker) (ITGAX/1243) , Biotin conjugate, 0.1mg/mL, Clone: [ITGAX/1243], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1243-500 |
| Hersteller Artikelnummer | BNCB1243-500 |
| Alternativnummer | BOT-BNCB1243-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human ITGAX protein fragment (aa 637-827) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a protein of 145 kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (M... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ITGAX/1243] |

| | |
|------------------------|--|
| Molekulargewicht | 145 kDa |
| UniProt | P20702 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |