

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD7 (T-Cell Leukemia Marker)(HuLy-m2), Biotin conjugate, 0.1mg/mL, Clone: [HuLy-m2], Mouse, Monoclonal BOT-BNCB1252-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD7 (T-Cell Leukemia Marker)(HuLy-m2), Biotin conjugate, 0.1mg/mL, Clone: [HuLy-m2], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNCB1252-100  |
| Hersteller Artikelnummer | BNCB1252-100  |
| Alternativnummer         | BOT-BNCB1252-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IF  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human thymocytes  |
| Konjugation              | Biotin  |
| Produktbeschreibung      | Recognizes a protein of 40 kDa, identified as CD7 (Workshop IV, Code T165). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of th... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [HuLy-m2]   |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 40 kDa  |
| UniProt                | <a href="#">P09564</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |