

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Oncoprotein(MYC275 + MYC909), Biotin conjugate, 0.1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal BOT-BNCB1269-500

Artikelname	c-Myc Oncoprotein(MYC275 + MYC909), Biotin conjugate, 0.1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1269-500
Hersteller Artikelnummer	BNCB1269-500
Alternativnummer	BOT-BNCB1269-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human c-myc protein
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a var...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MYC275 MYC909]

Molekulargewicht	62-64 kDa
UniProt	P01106
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user