

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 10 (Suprabasal Epithelial Marker) (DE-K10), Biotin conjugate, 0.1mg/mL, Clone: [DE-K10], Mouse, Monoclonal BOT-BNCB1273-100**

Artikelname	Cytokeratin 10 (Suprabasal Epithelial Marker) (DE-K10), Biotin conjugate, 0.1mg/mL, Clone: [DE-K10], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1273-100
Hersteller Artikelnummer	BNCB1273-100
Alternativnummer	BOT-BNCB1273-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Canine, Feline, Human
Immunogen	Cytoskeletal preparation extracted from human ectocervical epithelium
Konjugation	Biotin
Produktbeschreibung	This MAb recognizes a protein of 56.5 kDa, identified as cytokeratin 10 (CK10). CK10 is expressed in all suprabasal layers of the epidermis. In the epidermis, expression of CK10 strictly parallels the extent of differentiation, it is absent in the ba...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[DE-K10]
Molekulargewicht	56.5 kDa
UniProt	<a href="#">P13645</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user