

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Carcinoembryonic Antigen (CEA) / CD66 (C66/1292), Biotin conjugate, 0.1mg/mL, IgG2b, Clone: [C66/1292], Mouse, Monoclonal BOT-BNCB1292-100

| | |
|--------------------------|---|
| Artikelname | Carcinoembryonic Antigen (CEA) / CD66 (C66/1292), Biotin conjugate, 0.1mg/mL, IgG2b, Clone: [C66/1292], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1292-100 |
| Hersteller Artikelnummer | BNCB1292-100 |
| Alternativnummer | BOT-BNCB1292-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CEA protein |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes proteins of 80-200 kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MA... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [C66/1292] |

| | |
|------------------------|--|
| Molekulargewicht | 80-200 kDa |
| Isotyp | IgG2b |
| UniProt | P06731 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |