

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **HPV-16 E6 + HPV-18 E6 (Human Papilloma Virus 16/18)(HPV16/1295 + HPV18/1297), Biotin conjugate, 0.1mg/mL, Clone: [HPV16/1295 HPV18/1297], Mouse, Monoclonal BOT-BNCB1340-500**

|                          |   |
|--------------------------|---|
| Artikelname              | HPV-16 E6 + HPV-18 E6 (Human Papilloma Virus 16/18)(HPV16/1295 + HPV18/1297), Biotin conjugate, 0.1mg/mL, Clone: [HPV16/1295 HPV18/1297], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNCB1340-500  |
| Hersteller Artikelnummer | BNCB1340-500  |
| Alternativnummer         | BOT-BNCB1340-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Virus   |
| Immunogen                | HPV18 E6-beta-galactosidase fusion protein  |
| Konjugation              | Biotin  |
| Produktbeschreibung      | Human papilloma viruses (HPVs) can be classified as either high risk or low risk according to their association with cancer. HPV16 and HPV18 are the most common of the high risk group while HPV6 and HPV11 are among the low risk types. Approximately 9... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [HPV16/1295 HPV18/1297]   |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 16/17 kDa  |
| UniProt                | Not Applicable   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |