

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1342), Biotin conjugate, 0.1mg/mL, IgG2a, Clone: [CFTR/1342], Mouse, Monoclonal BOT-BNCB1342-100

| | |
|--------------------------|---|
| Artikelname | CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1342), Biotin conjugate, 0.1mg/mL, IgG2a, Clone: [CFTR/1342], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1342-100 |
| Hersteller Artikelnummer | BNCB1342-100 |
| Alternativnummer | BOT-BNCB1342-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human full-length CFTR |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structu... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [CFTR/1342] |
| Molekulargewicht | 165-170 kDa |
| Isotyp | IgG2a |
| UniProt | P13569 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |