

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8/18 (KRT8.18/1346), Biotin conjugate, 0.1mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal BOT-BNCB1346-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 8/18 (KRT8.18/1346), Biotin conjugate, 0.1mg/mL, Clone: [KRT8.18/1346], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1346-500 |
| Hersteller Artikelnummer | BNCB1346-500 |
| Alternativnummer | BOT-BNCB1346-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Cytoskeleton preparation from HeLa cells |
| Konjugation | Biotin |
| Produktbeschreibung | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb recognizes all simple epithelia including glandular epithelium, for example thyr... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KRT8.18/1346] |

| | |
|------------------------|---|
| Molekulargewicht | 52.5 kDa (CK8), 45 kDa (CK18) |
| UniProt | P05787 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |