

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Alpha-1-Antitrypsin (SERPINA1) (Hepatocellular & Histiocytic Marker) (AAT/1379), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [AAT/1379], Mouse, Monoclonal BOT-BNCB1379-500

| | |
|--------------------------|---|
| Artikelname | Alpha-1-Antitrypsin (SERPINA1) (Hepatocellular & Histiocytic Marker) (AAT/1379), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [AAT/1379], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1379-500 |
| Hersteller Artikelnummer | BNCB1379-500 |
| Alternativnummer | BOT-BNCB1379-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Alpha-1-Antitrypsin (AAT) protein fragment (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a protein of 54 kDa, which is identified as Alpha-1 antitrypsin (AAT). AAT is secreted and is a serine protease inhibitor whose targets include elastase, plasmin, thrombin, trypsin, chymotrypsin, and plasminogen activator. De... |
| Klonalität | Monoclonal |

| | |
|------------------------|---|
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [AAT/1379] |
| Molekulargewicht | 54 kDa |
| Isotyp | IgG1 |
| UniProt | P01009 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |