

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Lewis Y (Tumor Marker) (A70-A/A9), Biotin conjugate, 0.1mg/mL, Clone: [A70-A/A9], Mouse, Monoclonal BOT-BNCB1412-500**

Artikelname	Lewis Y (Tumor Marker) (A70-A/A9), Biotin conjugate, 0.1mg/mL, Clone: [A70-A/A9], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1412-500
Hersteller Artikelnummer	BNCB1412-500
Alternativnummer	BOT-BNCB1412-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Live Ls174T cells (human colon carcinoma cell line)
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a carbohydrate epitope present on tumor-associated Lewis Y antigen (Fucalpha1-2Galbeta1-4/3GlcNAcbeta). Lewis Y is expressed in large bowel tumors and colorectal carcinomas. It may be useful in the classification of human ren...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[A70-A/A9]

Molekulargewicht	Multiple
UniProt	Not Applicable
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user