

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Desmoglein-3 (Squamous Cell Marker) (5G11), Biotin conjugate, 0.1mg/mL, Clone: [5G11], Mouse, Monoclonal BOT-BNCB1439-500

| | |
|--------------------------|--|
| Artikelname | Desmoglein-3 (Squamous Cell Marker) (5G11), Biotin conjugate, 0.1mg/mL, Clone: [5G11], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1439-500 |
| Hersteller Artikelnummer | BNCB1439-500 |
| Alternativnummer | BOT-BNCB1439-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human DSG3 extracellular protein fragment (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a protein of 130 kDa, identified as Desmoglein-3 (DSG3). This MAbs is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein c... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [5G11] |
| Molekulargewicht | 130 kDa |
| UniProt | P32926 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |