

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Catenin, beta (p120) (5H10), Biotin conjugate, 0.1mg/mL, Clone: [5H10], Mouse, Monoclonal BOT-BNCB1444-500

| | |
|--------------------------|---|
| Artikelname | Catenin, beta (p120) (5H10), Biotin conjugate, 0.1mg/mL, Clone: [5H10], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1444-500 |
| Hersteller Artikelnummer | BNCB1444-500 |
| Alternativnummer | BOT-BNCB1444-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC, WB |
| Spezies Reaktivität | Gallus, Human, Mouse |
| Immunogen | Fusion protein consisting of the maltose binding protein fused to a 100 amino acid segment of the C-terminus of chicken beta-Catenin |
| Konjugation | Biotin |
| Produktbeschreibung | Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [5H10] |
| Molekulargewicht | 92 kDa |
| UniProt | P35222 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA pH 7.5-8.5 Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |