

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD3e (T-Cell Marker) (PC3/188A), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [PC3/188A], Mouse, Monoclonal BOT-BNCB1479-100

Artikelname	CD3e (T-Cell Marker) (PC3/188A), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [PC3/188A], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1479-100
Hersteller Artikelnummer	BNCB1479-100
Alternativnummer	BOT-BNCB1479-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Synthetic peptide corresponding to amino acids 156-168 (ERPPPVPNPDYEP) of the cytoplasmic domain of human CD3e
Konjugation	Biotin
Produktbeschreibung	The immunogen for this Mab is derived from the cytoplasmic domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closely ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[PC3/188A]
Molekulargewicht	20 kDa
Isotyp	IgG1
UniProt	P07766
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Not recommended for flow cytometry Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user