

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD45 / LCA (B-Cell Marker) (F10-89-4), Biotin conjugate, 0.1mg/mL, Clone: [F10-89-4], Mouse, Monoclonal BOT-BNCB1522-500**

Artikelname	CD45 / LCA (B-Cell Marker) (F10-89-4), Biotin conjugate, 0.1mg/mL, Clone: [F10-89-4], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1522-500
Hersteller Artikelnummer	BNCB1522-500
Alternativnummer	BOT-BNCB1522-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC
Spezies Reaktivität	Human
Immunogen	Human T-Lymphocytes
Konjugation	Biotin
Produktbeschreibung	CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[F10-89-4]
Molekulargewicht	180-220 kDa
UniProt	<a href="#">P08575</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user