

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD162 (Selectin P Ligand) (PSGL1/1601), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [PSGL1/1601], Mouse, Monoclonal BOT-BNCB1601-500**

Artikelname	CD162 (Selectin P Ligand) (PSGL1/1601), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [PSGL1/1601], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1601-500
Hersteller Artikelnummer	BNCB1601-500
Alternativnummer	BOT-BNCB1601-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length SELPLG (CD162) protein
Konjugation	Biotin
Produktbeschreibung	CD162 glycoprotein functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking duri...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[PSGL1/1601]

Molekulargewicht	43-120 kDa
Isotyp	IgG1
UniProt	<a href="#">Q14242</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user