

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CDX2 (GI Epithelial Marker) (CDX2/1690), Biotin conjugate, 0.1mg/mL, Clone: [CDX2/1690], Mouse, Monoclonal BOT-BNCB1690-500**

Artikelname	CDX2 (GI Epithelial Marker) (CDX2/1690), Biotin conjugate, 0.1mg/mL, Clone: [CDX2/1690], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1690-500
Hersteller Artikelnummer	BNCB1690-500
Alternativnummer	BOT-BNCB1690-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human CDX2 protein fragment (aa150-249) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CD...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[CDX2/1690]
Molekulargewicht	40 kDa
UniProt	<a href="#">Q99626</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL ELISA: 2-4 ug/mL for coating order Ab without BSA Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user