

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CTLA4 / CD152 (Negative Regulator of T-Cells)(L4P2F5.F10), Biotin conjugate, 0.1mg/mL, Clone: [L4P2F5.F10], Mouse, Monoclonal BOT-BNCB1696-500

Artikelname	CTLA4 / CD152 (Negative Regulator of T-Cells)(L4P2F5.F10), Biotin conjugate, 0.1mg/mL, Clone: [L4P2F5.F10], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1696-500
Hersteller Artikelnummer	BNCB1696-500
Alternativnummer	BOT-BNCB1696-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Bovine, Human, Mouse, Rat
Immunogen	A synthetic peptide from CTLA4 (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	CTLA4 (CD152) is a cell surface receptor that behaves as a negative regulator of the proliferation and the effector function of T cells. It contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[L4P2F5.F10]
Molekulargewicht	30-43 kDa

UniProt	P16410
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry