

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 15 (KRT15) (Basal Cell Marker) (KRT15/1699), Biotin conjugate, 0.1mg/mL, Clone: [KRT15/1699], Mouse, Monoclonal BOT-BNCB1699-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 15 (KRT15) (Basal Cell Marker) (KRT15/1699), Biotin conjugate, 0.1mg/mL, Clone: [KRT15/1699], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1699-500 |
| Hersteller Artikelnummer | BNCB1699-500 |
| Alternativnummer | BOT-BNCB1699-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Bovine, Human, Rat |
| Immunogen | Recombinant human KRT15 protein |
| Konjugation | Biotin |
| Produktbeschreibung | Keratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I keratin (keratins 9 to 23) and one type II keratin (keratins 1 to 8). Keratins demonstrate tissue and differentiation spec... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KRT15/1699] |

| | |
|------------------------|---|
| Molekulargewicht | 52 kDa |
| UniProt | P19012 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (frozen or formalin) 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |