

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) (IG1707R), Biotin conjugate, 0.1mg/mL, Clone: [IG1707R], Rabbit, Monoclonal BOT-BNCB1707-100**

Artikelname	Human IgG (Immunoglobulin Gamma Heavy Chain) (B-Cell Marker) (IG1707R), Biotin conjugate, 0.1mg/mL, Clone: [IG1707R], Rabbit, Monoclonal
Artikelnummer	BOT-BNCB1707-100
Hersteller Artikelnummer	BNCB1707-100
Alternativnummer	BOT-BNCB1707-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified human Ig Gamma Chain
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a protein of 75 kDa, identified as gamma heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[IG1707R]
Molekulargewicht	75 kDa
UniProt	<a href="#">P01861</a> , <a href="#">P01860</a> , <a href="#">P01859</a> , <a href="#">P01857</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user