

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Desmin (Muscle Cell Marker) (DES/1711), Biotin conjugate, 0.1mg/mL, Clone: [DES/1711], Mouse, Monoclonal BOT-BNCB1711-100**

Artikelname	Desmin (Muscle Cell Marker) (DES/1711), Biotin conjugate, 0.1mg/mL, Clone: [DES/1711], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1711-100
Hersteller Artikelnummer	BNCB1711-100
Alternativnummer	BOT-BNCB1711-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Gallus, Hamster, Human, Mouse, Rat
Immunogen	Recombinant human full-length desmin protein
Konjugation	Biotin
Produktbeschreibung	Cytoskeletal intermediate filaments (IFs) constitute a diverse group of proteins that are expressed in a highly tissue-specific manner. IFs are constructed from two-chain alpha-helicalcoiled-coil molecules arranged on an imperfect helical lattice, an...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[DES/1711]

Molekulargewicht	52 kDa
UniProt	<a href="#">P17661</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Immunofluorescence 1-2 ug/mL Flow cytometry 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user