

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Glycophorin A / CD235a (Erythrocyte Marker) (GYPA/1725R), Biotin conjugate, 0.1mg/mL, Clone: [GYPA/1725R], Rabbit, Monoclonal BOT-BNCB1725-500**

Artikelname	Glycophorin A / CD235a (Erythrocyte Marker) (GYPA/1725R), Biotin conjugate, 0.1mg/mL, Clone: [GYPA/1725R], Rabbit, Monoclonal
Artikelnummer	BOT-BNCB1725-500
Hersteller Artikelnummer	BNCB1725-500
Alternativnummer	BOT-BNCB1725-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length Glycophorin A protein
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 39 kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[GYPA/1725R]

Molekulargewicht	39 kDa
UniProt	<a href="#">P02724</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user