

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glycophorin A / CD235a (Erythrocyte Marker) (JC159), Biotin conjugate, 0.1mg/mL, Clone: [JC159], Mouse, Monoclonal BOT-BNCB1737-500

| | |
|--------------------------|---|
| Artikelname | Glycophorin A / CD235a (Erythrocyte Marker) (JC159), Biotin conjugate, 0.1mg/mL, Clone: [JC159], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1737-500 |
| Hersteller Artikelnummer | BNCB1737-500 |
| Alternativnummer | BOT-BNCB1737-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Formalin fixed membrane from a Hairy cell leukemia |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a sialoglycoprotein of 39 kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [JC159] |

| | |
|------------------------|--|
| Molekulargewicht | 39 kDa |
| UniProt | P02724 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |