

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD43 (T-Cell Marker) (SPN/1766R), Biotin conjugate, 0.1mg/mL, Clone: [SPN/1766R], Rabbit, Monoclonal BOT-BNCB1766-100

Artikelname	CD43 (T-Cell Marker) (SPN/1766R), Biotin conjugate, 0.1mg/mL, Clone: [SPN/1766R], Rabbit, Monoclonal
Artikelnummer	BOT-BNCB1766-100
Hersteller Artikelnummer	BNCB1766-100
Alternativnummer	BOT-BNCB1766-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human SPN protein
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[SPN/1766R]

Molekulargewicht	115 95 or 135 kDa
UniProt	P16150
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user