

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD43 (T-Cell Marker) (SPN/1766R), Biotin conjugate, 0.1mg/mL, Clone: [SPN/1766R], Rabbit, Monoclonal BOT-BNCB1766-500

| | |
|--------------------------|---|
| Artikelname | CD43 (T-Cell Marker) (SPN/1766R), Biotin conjugate, 0.1mg/mL, Clone: [SPN/1766R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNCB1766-500 |
| Hersteller Artikelnummer | BNCB1766-500 |
| Alternativnummer | BOT-BNCB1766-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human SPN protein |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [SPN/1766R] |

| | |
|------------------------|---|
| Molekulargewicht | 115 95 or 135 kDa |
| UniProt | P16150 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |