

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1775R), Biotin conjugate, 0.1mg/mL, Clone: [CFTR/1775R], Rabbit, Monoclonal BOT-BNCB1775-500**

Artikelname	CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1775R), Biotin conjugate, 0.1mg/mL, Clone: [CFTR/1775R], Rabbit, Monoclonal
Artikelnummer	BOT-BNCB1775-500
Hersteller Artikelnummer	BNCB1775-500
Alternativnummer	BOT-BNCB1775-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human CFTR fragment (aa258-385) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structu...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CFTR/1775R]
Molekulargewicht	165-170 kDa
UniProt	<a href="#">P13569</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user