BIOZOL
Diagnostica Vertrieb GmbH, Oehleckerring 11-13
22419 Hamburg, Germany
Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99
E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD8A (Cytotoxic- & Suppressor T-Cell Marker) (C8/1779R), Biotin conjugate, 0.1mg/mL, Clone: [C8/1779R], Rabbit, Monoclonal BOT-BNCB1779-100

Artikelname	CD8A (Cytotoxic- & Suppressor T-Cell Marker) (C8/1779R), Biotin conjugate, 0.1mg/mL, Clone: [C8/1779R], Rabbit, Monoclonal
Artikelnummer	BOT-BNCB1779-100
Hersteller Artikelnummer	BNCB1779-100
Alternativnummer	BOT-BNCB1779-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CD8a protein
Konjugation	Biotin
Produktbeschreibung	CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[C8/1779R]

Molekulargewicht	32 kDa
UniProt	P01732
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody|Immunofluorescence: 0.5-1 ug/mL|Immunohistology (formalin) 0.5 ug/mL|Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min|Optimal dilution for a specific application should be determined by user