

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD209 / DC-SIGN (Pathogen Receptor on Dendritic Cells) (C209/1781), Biotin conjugate, 0.1mg/mL, Clone: [C209/1781], Mouse, Monoclonal BOT-BNCB1781-100

| | |
|--------------------------|---|
| Artikelname | CD209 / DC-SIGN (Pathogen Receptor on Dendritic Cells) (C209/1781), Biotin conjugate, 0.1mg/mL, Clone: [C209/1781], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1781-100 |
| Hersteller Artikelnummer | BNCB1781-100 |
| Alternativnummer | BOT-BNCB1781-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human CD209 protein fragment |
| Konjugation | Biotin |
| Produktbeschreibung | DC-SIGN is a transmembrane receptor that is expressed on the surface of dendritic cells and macrophages. It is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses. The prot... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [C209/1781] |
| Molekulargewicht | 45 kDa |
| UniProt | Q9NNX6 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Immunohistology (formalin) 0.5 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |