

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD209 / DC-SIGN (Pathogen Receptor on Dendritic Cells) (C209/1781), Biotin conjugate, 0.1mg/mL, Clone: [C209/1781], Mouse, Monoclonal BOT-BNCB1781-500**

Artikelname	CD209 / DC-SIGN (Pathogen Receptor on Dendritic Cells) (C209/1781), Biotin conjugate, 0.1mg/mL, Clone: [C209/1781], Mouse, Monoclonal
Artikelnummer	BOT-BNCB1781-500
Hersteller Artikelnummer	BNCB1781-500
Alternativnummer	BOT-BNCB1781-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human CD209 protein fragment
Konjugation	Biotin
Produktbeschreibung	DC-SIGN is a transmembrane receptor that is expressed on the surface of dendritic cells and macrophages. It is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses. The prot...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[C209/1781]
Molekulargewicht	45 kDa
UniProt	<a href="#">Q9NNX6</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Immunohistology (formalin) 0.5 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user