

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Histone H1 (Nuclear Marker) (HH1/1784R), Biotin conjugate, 0.1mg/mL, Clone: [HH1/1784R], Rabbit, Monoclonal BOT-BNCB1784-500**

Artikelname	Histone H1 (Nuclear Marker) (HH1/1784R), Biotin conjugate, 0.1mg/mL, Clone: [HH1/1784R], Rabbit, Monoclonal
Artikelnummer	BOT-BNCB1784-500
Hersteller Artikelnummer	BNCB1784-500
Alternativnummer	BOT-BNCB1784-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant full-length human Histone H1 protein
Konjugation	Biotin
Produktbeschreibung	Eukaryotic histones are basic and water-soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed supealpha-helicalturn sequentially to form chromosomal fiber. Two molecules of each of...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[HH1/1784R]

Molekulargewicht	~30 kDa
UniProt	Multiple
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user