

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD40 / TNFRSF5 / CD40L-Receptor(T8P2G4*A6), Biotin conjugate, 0.1mg/mL, Clone: [T8P2G4*A6], Mouse, Monoclonal BOT-BNCB1802-500

| | |
|--------------------------|---|
| Artikelname | CD40 / TNFRSF5 / CD40L-Receptor(T8P2G4*A6), Biotin conjugate, 0.1mg/mL, Clone: [T8P2G4*A6], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB1802-500 |
| Hersteller Artikelnummer | BNCB1802-500 |
| Alternativnummer | BOT-BNCB1802-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Ovalbumin-conjugated synthetic peptide (RDPVCHPLGA) from the C-terminal of human CD40 |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody is specific for the soluble form of CD40. It is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin cl... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [T8P2G4*A6] |

| | |
|------------------------|--|
| Molekulargewicht | 43 kDa |
| UniProt | P25942 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |