

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TCL1 (T-Cell Marker) (TCL1/2078), Biotin conjugate, 0.1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal BOT-BNCB2078-100

| | |
|--------------------------|---|
| Artikelname | TCL1 (T-Cell Marker) (TCL1/2078), Biotin conjugate, 0.1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB2078-100 |
| Hersteller Artikelnummer | BNCB2078-100 |
| Alternativnummer | BOT-BNCB2078-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [TCL1/2078] |
| Molekulargewicht | 14 kDa |
| UniProt | P56279 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |