

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ATG5 (Autophagy Marker) (ATG5/2101), Biotin conjugate, 0.1mg/mL, Clone: [ATG5/2101], Mouse, Monoclonal BOT-BNCB2101-100

Artikelname	ATG5 (Autophagy Marker) (ATG5/2101), Biotin conjugate, 0.1mg/mL, Clone: [ATG5/2101], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2101-100
Hersteller Artikelnummer	BNCB2101-100
Alternativnummer	BOT-BNCB2101-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human ATG5 protein (around aa 1-119) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	Autophagy protein 5 (ATG5), in combination with autophagy protein 12, functions as an E1-like activating enzyme in a ubiquitin-like conjugating system. ATG5 is involved in several cellular processes, including autophagic vesicle formation, mitochondr...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[ATG5/2101]
Molekulargewicht	50-32 kDa
UniProt	Q9H1Y0
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: For coating use Ab at 1-5 ug/mL order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Western Blot 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user