

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2151), Biotin conjugate, 0.1mg/mL, Clone: [TACSTD2/2151], Mouse, Monoclonal BOT-BNCB2151-100

Artikelname	TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2151), Biotin conjugate, 0.1mg/mL, Clone: [TACSTD2/2151], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2151-100
Hersteller Artikelnummer	BNCB2151-100
Alternativnummer	BOT-BNCB2151-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[TACSTD2/2151]
Molekulargewicht	40 kDa
UniProt	P09758
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user