

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HCG-beta (Pregnancy & Choriocarcinoma Marker) (rHCGb/54), Biotin conjugate, 0.1mg/mL, Clone: [rHCGb/54], Mouse, Monoclonal BOT-BNCB2197-500

| | |
|--------------------------|---|
| Artikelname | HCG-beta (Pregnancy & Choriocarcinoma Marker) (rHCGb/54), Biotin conjugate, 0.1mg/mL, Clone: [rHCGb/54], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB2197-500 |
| Hersteller Artikelnummer | BNCB2197-500 |
| Alternativnummer | BOT-BNCB2197-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human hCG beta protein |
| Konjugation | Biotin |
| Produktbeschreibung | This MAb reacts with a protein of 22 kDa, identified as beta sub-unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [rHCGb/54] |

| | |
|------------------------|--|
| Molekulargewicht | 22 kDa |
| UniProt | P01233 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |