

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Kappa Light Chain (B-Cell Marker) (rL1C1), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [rL1C1], Mouse, Monoclonal BOT-BNCB2222-100

Artikelname	Human Kappa Light Chain (B-Cell Marker) (rL1C1), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [rL1C1], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2222-100
Hersteller Artikelnummer	BNCB2222-100
Alternativnummer	BOT-BNCB2222-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human B-Lymphoma Cells
Konjugation	Biotin
Produktbeschreibung	This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain, ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rL1C1]

Molekulargewicht	~22.5 kDa
Isotyp	IgG1
UniProt	P01601
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user