

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Insulin Receptor Alpha (INSR/2277R), Biotin conjugate, 0.1mg/mL, Clone: [INSR/2277R], Rabbit, Monoclonal BOT-BNCB2277-500

| | |
|--------------------------|--|
| Artikelname | Insulin Receptor Alpha (INSR/2277R), Biotin conjugate, 0.1mg/mL, Clone: [INSR/2277R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNCB2277-500 |
| Hersteller Artikelnummer | BNCB2277-500 |
| Alternativnummer | BOT-BNCB2277-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of extracellular domain of human Insulin Receptor alpha (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | The insulin receptor (INSR) is a heterodimeric protein complex that has an intracellular beta subunit and an extracellular alpha subunit, which is disulfide-linked to a transmembrane segment. The insulin ligand binds to the INSR and initiates molecu... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [INSR/2277R] |
| Molekulargewicht | 135 kDa |
| UniProt | P06213 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |